

Oral Pharmaceuticals in an Optometric Practice

Blair Lonsberry, MS, OD, MEd., FAAO
Professor of Optometry
Pacific University College of Optometry
blonsberry@pacificu.edu

Disclosures:

- Sun Pharmaceuticals: speakers bureau,
- Avellino: advisory board,
- Dompe: advisory board,
- RVL Pharmaceuticals: advisory board




1

2

Case

- 20 year old male presents with a red painful eye
 - Started that morning when he woke up
 - reports a watery discharge, no itching, and is not a contact lens wearer
- SLE:
 - See attached image with NaFl stain



3

Herpes Simplex Virus (HSV) Keratitis: Clinical Features

- Characterized by primary outbreak and subsequent reactivation
 - Primary outbreak is typically mild or subclinical (90% of people are asymptomatic)
 - Most clinical ocular infections are manifestations of virus reactivation; ocular involvement occurs in fewer than 5% of primary infections
- After primary infection, the virus becomes latent in the trigeminal ganglion or cornea
 - The majority of ophthalmic HSV cases are unilateral, with recurrences affecting the same eye. Bilateral disease (not necessarily concurrent) occurs in 1-12% of cases and is more common in patients with atopy or other immune abnormalities
- Stress, UV radiation, and hormonal changes can reactivate the virus
- Lesions are common in the immunocompromised (i.e. recent organ transplant or HIV patients)



3

4


Herpes Simplex Virus Keratitis

- **Epithelial Keratitis:**
 - Symptoms:
 - Ocular irritation, redness, photophobia, watering, blurred vision
 - Signs:
 - Swollen opaque epithelial cells arranged in a coarse punctate or stellate pattern
 - Central desquamation results in a dendrite***
 1. Central ulceration
 2. Terminal end bulbs
- ***Corneal sensation is reduced***



5


Dendritic Ulcers



6

6

HSV Geographic Ulcer



https://www.researchgate.net/figure/Geographic-corneal-ulcer-caused-by-herpes-simplex-virus-keratitis_fig1_267320111



7

Pediatric HSV Keratitis

- pediatric herpes simplex keratitis has an 80% risk of recurrence, a 75% risk of stromal disease, and a 30% rate of misdiagnosis
- 80% of children with herpes simplex keratitis develop scarring, mostly in the central cornea
 - results in the development of astigmatism
 - 25% of children have more than 2 D of astigmatism, most of which is irregular
- consider pediatric HSV when a patient has unilateral recurrent disease in the anterior segment



8


Herpes Simplex Virus Keratitis Management

- Topical:
 - Viroptic (trifluridine) q 2h until epi healed then taper down for 10-14 days.
 - Viroptic is toxic to the cornea.
 - Zirgan (ganciclovir) available, use 5 times a day until epi healed then 3 times for a week (US only)



9

First pass metabolism



10

10

Anti-Viral Medication

Drug	Mechanism of Action	Bioavailability	Dosing	Side Effects
Acyclovir	Acyclovir interferes with DNA synthesis inhibiting viral replication	10-30% gets absorbed Short ½ life *Metabolized in kidneys	Simplex: 400 mg 5x/day Zoster: 800 mg 5x/day	Overall very safe Nausea, vomiting, headaches, dizziness, confusion
Valacyclovir	Acyclovir pro-drug Equivalent to acyclovir but better for pain management	95% converted to acyclovir* Better bioavailability and longer 1/2 life	Simplex: 500 mg tid Zoster: 1 g tid	Same as acyclovir
Famciclovir	Inhibits DNA chain elongation It is metabolized to penciclovir where it is active 10-20x as long as acyclovir	Superior to acyclovir*	Simplex: 250 mgTID Zoster: 500 mg TID	Same as acyclovir

11

HSV Stromal Disease



- HSV Stromal disease is an immune-mediated disease
 - Stromal involvement is rarely an initial ocular finding, accounting for fewer than 2% of initial presentations but for 20 – 60% of recurrent corneal disease
- Increased risk of scarring and high risk of poor visual prognosis
- Requires corticosteroids (HEDS: corticosteroid reduced risk of progression by 68%)
 - Without epithelial defect: corticosteroids and prophylactic anti-viral dosage
 - With epithelial defect: active infection anti-viral dosage with judicious corticosteroids



12

How much to dose steroid?

- HEDS used QID of *prednisolone phosphate*
- Current Recommendations:
 - Mod – severe (especially with neo): 1% Prednisolone or Lotemax QID to 6x/day
 - Want the lowest dose needed to control the inflammation
 - AAO EBM Treatment Guideline 2014
 - Topical steroid for 10 weeks (this is based on HEDS results) with oral antiviral



13

HSV Epithelial Keratitis

- Treatment Regimen:
 - Oral Valtrex 500 mg 3x/day for 7-10 days
 - Artificial tears
 - L-Lysine 2 grams daily?
 - Proven to “slow down” and retard the growth of the herpes virus and inhibit viral replication
 - Debride the ulcer?
 - Prior to topical antiviral therapy debridement was treatment of choice
 - Generally try to avoid use of sharp instruments and use of cotton swab and anesthetic
- RTC 1 day, 4 days, 7 days



14

Herpes Simplex Keratitis

- Prophylactic Treatment:
 - Reduces the rate of recurrence of epithelial and stromal keratitis by ≈ 50%
 - Acyclovir 400 mg BID
 - Valtrex 500 mg QD
 - Famvir 250 mg QD
 - L-lysine 1 gram/day:
 - Proven to “slow down” and retard the growth of the herpes virus and inhibit viral replication
- Frequent debilitating recurrences, bilateral involvement, or HSV infection in a monocular patient



15


Prophylaxis??

• Pitfalls to Prophylaxis:

- Reduction of recurrence does not persist once drug stopped
- Resistance????
 - van Velzen, et. al., (2013) demonstrated that long-term ACV prophylaxis predisposes to ACV-refractory disease due to the emergence of corneal ACVR HSV-1.



16



17

Oral Pain Management

18

Pain Management: Oral Analgesics

- Conditions potentially requiring use of oral analgesics:
 - Corneal ulcers
 - Herpes simplex/zoster
 - Post-surgical
 - Trauma
 - Thermal burns
 - Periorbital infections (e.g. dacryocystitis/preseptal)



19

Oral Analgesics: Guidelines

- Make the proper diagnosis first (ie. Don't prescribe without knowing what you are prescribing for!)
- Treat the underlying cause for the pain
- **treat the pain at presentation..don't wait!**
- **treat pain continuously over a 24 hour schedule**
- Non-prescription drugs should be first choice and tend to be low cost
- Treat patients with the simplest and safest means to alleviate pain



20

Oral Analgesics: Guidelines

- **Mild to moderate pain is often successfully treated with acetaminophen or NSAIDs**
- **Moderate to severe pain is best treated with opioid analgesics**
- Adjunctive treatments are very valuable in pain management:
 - Mydriatic/cycloplegic useful for ocular pain
 - Bandage CL or pressure patch
 - Topical skin: Zostrix Cream (Capsaicin), EMLA Cream (lidocaine 2.5% and prilocaine 2.5%)



21

Systemic NSAID's

- **NSAID's are the drug of choice for treating mild to moderate ocular pain.**
 - Very beneficial for treating systemic inflammation as well.
- All NSAID's are rapidly absorbed from the GI tract, highly bound in the plasma, and capable of crossing the blood-brain barrier.
- **Exhibit a "ceiling effect" – there is a dosage beyond which no further analgesia occurs.**
 - Produce no tolerance or dependence, increasing their safety profile.
- **Variability exists in patient responses to NSAID's**
 - No definitive recommendation on treatment can be given.
 - If one NSAID does not work – TRY ANOTHER.



22

Aspirin (ASA)

- Weak organic acid.
- Oldest non-opioid analgesic available today.
- Very good anti-inflammatory and antipyretic properties.
 - Adult Dosage: 325 – 650 mg every 4 hours
 - Do not exceed 4 grams/day
 - **Most Common use of ASA:** Inhibit platelet aggregation in patients with history of heart attacks and heart surgery.
 - Most common dosing is **81 mg/day**
- **Largely replaced as treatment for pain associated with inflammation by the other classes of NSAID's due to the frequent side effects**
 - GI Distress: Inhibit prostaglandin synthesis and the production of a mucous lining on the stomach leading to increased gastric acid secretion.



23

2019 Aspirin Recommendations

- American College of Cardiology:
 - **Aspirin should be used infrequently in the routine primary prevention of ASCVD because of lack of net benefit.**
 - low-dose aspirin should not be routinely given as a preventive measure to adults 70 years and older or to any adult who has an increased risk of bleeding
 - Just updated (2021): no longer recommended for patients 60 and older, and for patients aged 40-59 it should be on a case to case basis.



24

Ibuprofen

- Mild to moderate pain, fever
- Adult analgesic dose: 200-400mg q4-6 hours
 - Maximum Dosage: 2400 mg/day for pain (approved for 3200 mg/day in arthritis treatment)
- OTC: 200 mg tabs (US) 400 mg and 600 mg (Canada)
- Rx: 300, 400, 600, 800mg tabs
 - Can prescribe 800 mg q8hrs
- Peak levels 1-2 hours
- Most renal toxic of all the NSAID's
- Brand Names: Motrin, Advil, and Nuprin



25

Naproxen Sodium

- OTC: 220 mg (Aleve®)
- Rx: 550 mg tablets (Anaprox® and Crysanal®)
- For mild to moderate pain
- Adult Dose:
 - OTC: 2 tablets first dose, then 1 tablet 8-12 hours (max dose 1250 mg)
 - Rx: 550 initial dose, followed by 275 (half tablet) every 6-8 hours.
 - Maximum Dose: 1375mg/day.



26

Indomethacin (Indocin)

- used to treat moderate to severe osteoarthritis, rheumatoid arthritis, gouty arthritis, or ankylosing spondylitis.
- Usual Adult Dosage for Pain: 25-50 mg two to three times/day
- Rx Only: 25, 50 and 75mg capsules
- Mainly used as a short-term anti-inflammatory especially for conditions that do not respond to less toxic NSAIDs.
 - Indomethacin has a very high level of intolerance compared to other NSAID's.
- Oral NSAID most widely used in Tx of ocular inflammation
 - E.g. Scleritis treatment 75 mg BID



27

Cox-2 Inhibitors

- Selective agents for only COX-2 designed to protect the GI system from the side effects seen with NSAID's
- It is approved for the management of the signs and symptoms of osteoarthritis, rheumatoid arthritis, JRA (in patients >2), ankylosing spondylitis and acute pain
- Major agent available on the market is Celecoxib (Celebrex).
 - Other agents Valdecoxib (Bextra) and Rofecoxib (Vioxx) were removed from the market due to increased risk of heart attacks and strokes.
- Available: 50, 100, 200 and 400 mg capsules
- Osteoarthritis Dosage: 100 mg BID or 200 mg single dose daily
- RA: 100 to 200 mg BID daily



28

Contraindications to NSAIDs

- Avoid in:
 - Pregnancy (especially the late trimesters)
 - Active Peptic Ulcer Disease
 - Cross Sensitivity to ASA
 - Previous Hypersensitivity to NSAIDs
 - Chronic Renal Insufficiency
- At Risk Patients Include:
 - Dehydration
 - HTN or CHF
 - Use of ACE Inhibitors, diuretics and B-blockers
 - Higher doses of NSAIDs and chronic therapy extending beyond a week will be more likely to increase BP
 - Advanced Age



29

NSAIDs Black Box Warning

- BLACK BOX WARNING:
 - May increase the risk of serious thrombotic events, MI, and stroke.
 - Increase risk of serious GI adverse effects such as bleeding, ulcer, and perforation.



30

NSAID-related ulcers

- COX-2 inhibitors such as celecoxib (Celebrex) are less likely to cause ulcers than aspirin
- Proton pump inhibitors (e.g. Losec[®], Prevacid[®] or Prilosec[®]) help to offset the risk of NSAID-related stomach ulcers
 - patients should be treated with concomitant proton pump inhibitors once daily, which results in ulcer healing rates of approximately 80% at 8 weeks in patients continuing to take NSAIDs



31

Acetaminophen



- Mechanism of Action is not well understood.
 - Possibly some CNS component
 - Very weak inhibitor of prostaglandin synthesis
- One of the most commonly used analgesics for mild to moderate pain
 - Equal analgesic properties to ASA unless associated with inflammation, where it is less effective.

Take home: Good for pain; Good for fever;
No effect on inflammation



32

Acetaminophen



- Typical Adult Dosage (FDA Based):
 - 650 mg every 4 - 6 hours for Regular Strength (2 X 325)
 - Cannot take more than 10 caplets in 24 hours.
 - 1000 mg every 6 hours for Extra Strength (2 X 500)
 - Cannot take more than 6 caplets in 24 hours.
 - 1300 mg every 8 hours for Extended Release (2 X 650)
 - Cannot take more than 6 capsules in 24 hours.
- Daily dose of acetaminophen should not exceed 3 grams
 - This has been recently changed from 4000 mg which can be done with doctor approval.
- Should only be used for short term therapy
- Exhibits a ceiling effect, like NSAIDs and ASA



33

Dangers of Acetaminophen

- Acetaminophen overdose is the leading cause of liver failure in the U.S.
 - It sends 56,000 people to the emergency room annually and causes approximately 400 deaths yearly.
- Acetaminophen is used in so many products, people are often unaware that they are taking it, leading to more overdoses.
 - Combined with agents to get wide range of symptom coverage.
 - Antihistamines such as diphenhydramine – Tylenol PM
 - Diuretics such as Pyrilamine maleate – Midol Complete
 - Cough Suppressants such as Dextromethorphan – Nyquil



34

Consider Combining APAP with NSAID's for Mild to Moderate Pain Relief

1:00 pm: Two 325mg Tylenol
3:00 pm: Two 200mg Ibuprofen
5:00 pm: Two 325mg Tylenol
7:00 pm: Two 200mg Ibuprofen

Alternated every 2 hours while awake
• Each medication is q 4 hours.



35

Oral Analgesics: Guidelines

- Never exceed maximum recommended dosages:
 - ASA: 4 grams/day
 - Acetaminophen: 4 grams/day???? (newer data suggest should be closer to 3.2 grams/day)
 - Ibuprofen: 2400 mg/day OTC and up to 3200 mg/day prescription (for RA)
 - Naproxen: 1250/day
 - Naproxen sodium: 1375/day
 - Codeine: 360 mg/day



36

Gabapentin (Neurontin^R)

- Classified as an anticonvulsant drug
- Additionally, used in the treatment of patients with chronic pain
- Gabapentin, is not currently classified as a controlled substance in most states, however, its abuse potential is still being investigated.
 - Kentucky, Michigan, Tennessee, West Virginia ??, Virginia ??, and Ohio ?? have reclassified gabapentin as a Schedule V controlled substance.



37

Gabapentin (Neurontin^R)

- Gabapentin has primarily been studied and found effective for the treatment of postherpetic neuralgia and painful diabetic neuropathy; evidence for efficacy in other types of neuropathic pain is limited
- Treatment with gabapentin should be initiated at a low dose with gradual increases until pain relief or dose-limiting adverse effects are achieved.
- Dosage:
 - Day 1 single 300 mg dose
 - Day 2 600 mg dose
 - Day 3 900 mg dose
 - Can be titrated up all the way to 1800 mg/day



38

Opioids vs Ibuprofen + Acetaminophen

- 1000mg acetaminophen and 400 mg ibuprofen
- Ibuprofen Plus Acetaminophen Equals Opioid Plus Acetaminophen for Acute Severe Extremity Pain. Am Fam Physician. 2018;97(5):348
- Effect of a Single Dose of Oral Opioid and Nonopioid Analgesics on Acute Extremity Pain in the Emergency Department: A Randomized Clinical Trial. JAMA. 2017;318(17):1661-1667.
 - no statistically significant or clinically important differences in pain reduction at 2 hours among single-dose treatment with ibuprofen and acetaminophen or with 3 different opioid and acetaminophen combination analgesics



39

Opioids Side Effects

- Side Effects are very hard to predict because opioids can cause CNS depression or stimulation.
- CNS Side Effects
 - Dizziness, lightheadedness, sedation, and drowsiness are the most common.
 - Mood elevation (euphoria) and disorientation can occur in some patients.
 - Exacerbated if used in combination with alcohol, depression medications such as tricyclic antidepressants, anticholinergics, antihistamines, anti-seizure medications, or muscle relaxants, etc.
 - Visual symptoms such as blurry vision, miosis, and diplopia can occur.



41

Opioid Side Effects

- Nausea and Vomiting (more common in ambulatory pts.)
- Constipation
 - Can be relieved by OTC docusate sodium (Colace).
- Respiratory Side Effects
 - Respiratory Depression
 - Most serious side effect of the opioids



42

Patient Education

- Avoid all depressants – especially using along with alcohol.
- Must educate all patients of risks of these symptoms and caution them for driving or operating dangerous machines.
- Stomach upset can be helped by consuming the medication with food.
- Watch for signs of breathing difficulty or changes in blood pressure.



43

Scheduled Medications – Most Opioids

Schedule	Description	Optometric Medications
I	Not commercially available; no approved indication	
II	Very addictive medications that are accepted for medicinal use	Oxycodone = OxyContin, OxyFast Oxycodone + APAP = Percocet Hydromorphone (Dilaudid) Codeine Sulfate = Codeine Generic Meperidine (Demerol) Hydrocodone + APAP = Lortab Hydrocodone + Ibuprofen = Generic
III	Significant abuse risk, but less potent than I or II. May still contain narcotics.	Codeine + APAP = Tylenol 3 and Tylenol 4
IV	Relatively low abuse potential and limited risk	Tramadol
V	Very limited abuse potential. May be OTC in some states.	Cough medicine with codeine

44

Schedule II Opioids: Hydrocodone

- Approximately 6X more potent than codeine
- Milder Side Effects than Codeine: Less constipation and sedation.
- Clinically believed to cause more euphoria than codeine, but this is not backed by clinical studies.



46

Opioids: Codeine

- Analgesic effect occurs within 20 minutes of ingestion and reaches a maximum at 1 – 2 hours
- Ceiling effect occurs.
- Usually administered in combination with acetaminophen .
 - Tylenol 1 (222): codeine 8 mg, 300 mg acetaminophen and 15 mg caffeine (Canada)
 - Tylenol 3 = Codeine 30 mg and Acetaminophen 300 mg
 - Dosage: 1-2 tablets every 4 hours.
 - Tylenol 4 = Codeine 60 mg and Acetaminophen 300 mg
 - Dosage: 1 tablet every 4 – 6 hours
- Also available as generic with 15, 30, or 60 mg of Codeine with 300 mg of Acet. or elixer of 12 mg codeine + 120 mg Acet. per 5 mL.
 - Elixer can be used in children for pain management if >3 years.



45

Opioids: Hydrocodone

- Used in combination with APAP and Ibuprofen.
 - Lortab: Hydrocodone 5, 7.5, and 10 mg with APAP 325 mg
 - Dosage: 1-2 tablet every 4-6 hours
 - Lortab Elixer: Hydrocodone 10 mg with APAP 300 / 15 mL
 - Dosage: 3 tsp every 4-6 hours
- Generic:
 - Hydrocodone bitartrate 5, 7.5, 10 mg and acetaminophen 300 mg
 - Hydrocodone bitartrate 5, 7.5, 10 mg and acetaminophen 325 mg
- Generic Elixer:
 - Hydrocodone bitartrate 7.5, 10 mg and acetaminophen 325 mg per 15 mL
- Generic:
 - Hydrocodone bitartrate 5, 7.5, 10 mg and ibuprofen 200 mg



47

Schedule II Opioids: Oxycodone

- Approximately 10-12X more potent than codeine
 - As potent as parenteral morphine when given orally.
- Lower level of side effects in comparison to morphine, but high level of euphoria produced, thus higher level of abuse risk.



48

Opioids: Oxycodone

- Available in combination with APAP (combinations with ASA or Ibuprofen discontinued in US).
- Dosage: 1 tablet every 4-6 hours
 - Endocet tablets:
 - 2.5, 5, 7.5 or 10 mg Oxycodone with 325 mg Acetaminophen
 - Nalocet:
 - Oxycodone hydrochloride 2.5 mg and acetaminophen 300 mg
 - Percocet Tablets
 - 2.5, 5, 7.5 or 10 mg Oxycodone with 325 mg Acetaminophen
 - Prolate:
 - 5, 7.5 or 10 mg Oxycodone with 300 mg Acetaminophen
 - Generic:
 - 2.5, 5, 7.5 or 10 mg Oxycodone with 325 mg Acetaminophen



49

Schedule IV: Tramadol

- Central acting narcotic
 - Synthetic analogue of codeine.
 - Binds to mu receptors and inhibits norepinephrine and serotonin reuptake.
 - Potential for abuse is very low, but has occurred.
- Generic: 50 mg, 100 mg
- **Dosage: 50 – 100 mg q4 – 6 hours.**
 - Analgesia occurs after 1 hour.
 - Maximum dose: 400 mg/day



50

Tramadol Extended Release

- ConZip:
 - Available dosages of 100, 200, and 300 mg extended.
 - Begin taking 100 mg daily X 5 days
 - Increase by 100 mg if relief not met to 200 mg X 5 days.
 - 300 mg maximum daily.
- Generic: 100 mg, 200 mg, 300 mg
- Does not work on all patients – some need heavy doses every 4-6 hours.
- More for chronic pain control.



51

Tramadol + APAP


- **Generic: Combination of:**
 - 325 mg of APAP
 - 37.5 mg of Tramadol
- **Dosage: 2 tablets every 4 – 6 hours**
- Max: 8 tablets daily



52

Preseptal Cellulitis

- Infection and inflammation located anterior to the orbital septum and limited to the superficial periocular tissues and eyelids.
- **Usually follows sinus infection or internal hordeolum** (possibly trauma)
- Eyelid swelling, redness, ptosis, pain and low grade fever.



53

Differentiating Orbital vs. Preseptal

FINDING	ORBITAL	PRESEPTAL
Visual Acuity	Decreased	Normal
Proptosis	Marked	Absent
Chemosis and Hyperemia	Marked	Rare/Mild
Pupils	RAPD	Normal
Pain and Motility	Restricted and Painful	Normal
IOP		Normal
Temperature	102 - 104	Normal/mild elevation
HA and Assoc. Symptoms	Common	Absent


Treatment: Orals for Preseptal, Often IV for Orbital



54

Preseptal Cellulitis

- Tx:
 - **Clavulin (Augmentin) 500 mg TID or 875 mg BID for 5-7 days**
 - **Keflex 500 mg QID 5-7 days**
 - or if moderate to severe IV Fortaz (ceftriaxone) 1-2 g q8h.
 - If MRSA possible, consider Bactrim/Septa



55

Antibiotic Resistance

- Microorganism that was originally in the spectrum of activity is no longer susceptible to the drug.
- **Mechanisms of Resistance Include:**
 - Producing an enzyme capable of destroying or inactivating the antibiotic.
 - Altering the target site receptor for the antibiotic so as to reduce or block its binding.
 - Preventing the entry of the antibiotic into the bacterial cell or actively transporting the antibiotic out.



56

Evolution of Bacterial Resistance

- <https://www.youtube.com/watch?v=plVk4NVIUh8>



57

Avoiding Resistance

- Bacterial resistance is a natural result of mutation.
- **Antibiotics cause a faster rate of selection against these resistant bacteria if not prescribed correctly.**
 - Avoid prescribing for non-bacterial infections.
 - Avoid sublethal doses (attack to kill all).
 - Avoid intermittent use.
 - Always complete the full dosage for an appropriate length of time.
 - NEVER TAPER AN ANTIBIOTIC below recommended dosing schedule!



58

Preventing Resistance

- **The IDSA suggests 5-7 days is long enough to treat a bacterial infection** without encouraging resistance in adults, though children should still get the longer course
- this is different than previous guidelines of treating infections from 10-14 days.



3/14/2023

59

MRSA

- **Healthcare-associated methicillin-resistant *Staphylococcus aureus* (HA-MRSA)** is associated with severe, invasive disease in hospitalized patients
- **Community-associated methicillin-resistant *S. aureus* (CA-MRSA)** is most often associated with skin and soft tissue infections in young, healthy individuals with no recent healthcare exposure



60

Consider Covering for MRSA

1. Hx of non-response to amoxicillin or Augmentin
2. Hx of previous MRSA infections
3. Infection did not start at lid margin like a regular hordeolum but more superior like near the eyebrow area
4. Hx of recent incarceration or hospitalization or in nursing home
5. health care worker
6. pain outside clinical presentation



61

61

Antibiotic Associated Diarrhea (AAD)

- The most common side effects of antibiotics are gastro-intestinal, such as nausea and diarrhea
- AAD arises when the antibiotic disrupts the ecology of the intestinal microbiota, by altering the diversity and numbers of bacteria in the gut.
- Diarrhea is most frequently associated with the use of broad-spectrum antibiotics (e.g. amoxicillin)

Agarwala, V., Kivi, C.A.M., Rijers, G. et al. A practical guide for probiotics applied to the case of antibiotic-associated diarrhea in The Netherlands. BMC Gastroenterol 18, 103 (2018)



62

AAD and Probiotics

- The core benefit of probiotics is exercised by contributing to the maintenance of a balanced microbiota and therefore by creating a favorable gut environment
- The efficacy of probiotics in preventing AAD depends on the dose.
 - A daily intake of at least 5×10^9 CFU is associated with significant efficacy for AAD and it has been shown that higher probiotic dose is linked to greater efficacy
 - Example: The probiotic content of yogurt products can range from 90 to 500 billion CFU per serving

Agarwala, V., Kivi, C.A.M., Rijers, G. et al. A practical guide for probiotics applied to the case of antibiotic-associated diarrhea in The Netherlands. BMC Gastroenterol 18, 103 (2018)



63

5 Facts About Penicillin Allergy

- Approximately 10% of all U.S. patients report having an allergic reaction to a penicillin class antibiotic in their past.
 - When evaluated, fewer than 1% of the population are truly allergic to penicillins
- Approximately 80% of patients with IgE-mediated penicillin allergy lose their sensitivity after 10 years
- Broad-spectrum antibiotics are often used as an alternative to penicillins. The use of broad-spectrum antibiotics in patients labeled "penicillin-allergic" is associated with higher healthcare costs, increased risk for antibiotic resistance, and suboptimal antibiotic therapy.
- Correctly identifying those who are not truly penicillin-allergic can decrease unnecessary use of broad-spectrum antibiotics.

<https://www.cdc.gov/antibiotic-use/community/pdf/penicillin-factsheet.pdf>



64

Penicillins: Clavulin (Augmentin)

- Clavulin (Augmentin) is amoxicillin with potassium clavulanate (clavulanic acid 125 mg).
- Clavulanate is a B-Lactamase inhibitor which reduces a bacteria's ability to negate the effect of the amoxicillin by inactivating penicillinase (enzyme that inactivates the antibiotic effect).
 - Dicloxacillin can also be used in infections due to penicillinase-producing staph.



65

Penicillins: Clavulin (Augmentin)

- Clavulin (Augmentin) is very effective for skin and skin structure infections such as:
 - dacryocystitis,
 - internal hordeola,
 - preseptal cellulitis.
- Treatment of:
 - otitis media,
 - sinusitis,
 - lower respiratory and urinary infections.
- Given prophylactically to dental surgery patients.



66

Penicillins: Clavulin (Augmentin)

- It has low:
 - GI upset,
 - allergic reaction and anaphylaxis.
- Serious complications include:
 - anemia,
 - pseudomembranous colitis and
 - Stevens-Johnson syndrome.



67

Penicillins: Augmentin.

Adults:

- **250 TID, 500 mg tab BID-TID depending on what you are treating (also available in chewable tablets and suspension)**
- **or 875 mg q 12hr (bid)**

- 1000 mg XR: q12 hr and not for use in children <16

Peds: <3 mos 30mg/kg/day divided q12hrs using suspension

- >3 mos 45-90mg/kg/day divided q12hrs (otitis media 90mg for 10 days)



68

Cephalosporins

- **Closely related structurally and functionally to the penicillins**
 - **have the same mode of action,**
 - affected by the same resistance mechanisms.
 - tend to be more resistant to B-lactamases.
- **Classified as 1st, 2nd, 3rd, 4th and now 5th generation** based largely on their bacterial susceptibility patterns and resistance to B-lactamases.
- Typically administered IV or IM, **poor oral absorption.**



69

- 1st generation: **cefadroxil (Duricef)**, **cefazolin (Ancef)**, **cephalexin (Keflex)**, and cephalothin
- 2nd generations: **cefaclor (Ceclor)**, cefprozil, **cefuroxime (Zinacef)**, cefotetan, cefoxitin
- 3rd generation: cefdinir (**Omnicef**), cefixime, **cefotaxime (Claforan)**, **cefazidime (Fortaz)**, ceftibuten, ceftizoxime, **ceftriaxone (Rocephin IM/IV)**.
- 4th generation: cefepime
- **Keflex, Ceclor, Omnicef**, (all orally administered) are effective against most gram positive pathogens and especially good for skin and soft tissue infections.



71

• Keflex (cephalexin):

- treatment of respiratory, GI, skin and skin structure, and bone infections as well as otitis media
- Adults: 250-1000 mg every 6 hours
 - **typical dosing 500 every 6 hours**
- Children: 25-100 mg/kg/day divided 6-8 hours



72

Cephalosporins

- Cefaclor (Ceclor) (2nd generation):
 - Immediate-release: 250 to 500 mg every 8 hours
 - Extended-release: 500 mg every 12 hours

Note: An extended-release tablet dose of 500 mg twice daily is clinically equivalent to an immediate-release capsule dose of 250 mg 3 times daily; an extended-release tablet dose of 500 mg twice daily is **NOT** clinically equivalent to 500 mg 3 times daily of other cefaclor formulations.



73

Cephalosporins

- **Cefdinir (Omnicef no longer available in the US, available as generic)**
 - Used in the treatment of community acquired pneumonia, acute flare ups of chronic bronchitis, acute maxillary sinusitis and tonsillitis.
- Adult dosing:
 - comes in 300 mg capsules and recommended dosing is 600 mg per day (single dose is equivalent to 300 every 12 hours)



74

Co-Trimoxazole (Bactrim/Septra)

- Combination of trimethoprim and sulfamethoxazole (TMX-SMX)
 - shows greater antimicrobial activity than equivalent quantities of either drug alone.
- Has broader spectrum of action than the sulfa's and is effective in treating:
 - UTIs and respiratory tract infections
 - often considered for treatment of MRSA skin infections



75

75

Co-Trimoxazole (Bactrim/Septra)

- Available:
 - Bactrim/Septra tablets SS (standard strength):**
 - contains 80 mg trimethoprim and 400 mg sulfamethoxazole
 - dosing **2 tablets every 12 hours**
 - Bactrim DS/Septra DS (Double Strength)**
 - contains 160 mg trimethoprim and 800 mg sulfamethoxazole
 - Dosing **1 tablet every 12 hours**



76

Co-Trimoxazole (Bactrim/Septra)

- Contraindication!!!!**
 - Methotrexate (MTX)** is a folic acid antagonist used for the treatment of many autoimmune diseases (e.g. rheumatoid arthritis, JIA, lupus etc)
 - TMX-SMX is an inhibitor of folic acid metabolism and can cause **bone marrow suppression**.
 - TMX-SMX is also known to decrease the renal excretion of MTX.
 - When used in combination, the potential for toxicity is substantial.
- Drink plenty of fluids to prevent kidney stones.
 - Interacts with diuretics and ACE inhibitors
- Exposure to sunlight, even for brief periods of time, may cause severe sunburn or skin rash, redness, itching, or discoloration.



Copyrights apply

77

Oral antimicrobial therapy for treatment of skin and soft tissue infections due to methicillin-resistant *Staphylococcus aureus* (MRSA) in adults

Treatment	Adult dose
Preferred agents*	
Trimethoprim-sulfamethoxazole (cotrimoxazole)	1 or 2 DS tablets twice daily [†]
Clindamycin	450 mg orally 3 times daily
Doxycycline	100 mg orally twice daily
Minocycline	200 mg orally once, then 100 mg orally twice daily
Alternative agents[‡]	
Linezolid	600 mg orally twice daily
Tedizolid	200 mg orally once daily
Delafloxacin	450 mg orally twice daily
Omadacycline	300 mg orally once daily

From Up to Date: MRSA in adults, accessed 09/06/2022

78


Case Example

- 67 YOF
- HA and vision loss x 2 days
- OHx: unremarkable
- LEE: 3 days ago!
- MHx: unremarkable


Case courtesy of Dr. Tammy Thien



79



80



Case courtesy of Dr. Tammy Thar



Pacific University
Oregon



81

Minocycline?

- Proposed mechanisms
 - MMPs (MMP-9)**
 - Increase in MMP-9 disrupt blood brain barrier and are linked to poor functional recovery
 - Anti-inflammatory**
 - Reduction in microglial activation**
 - microglial activation is believed to play a central role in neuroinflammation and pathological progression of ischemic tissue
 - Nitric oxide (NO) production**
 - NO plays a neuroprotective role in acute ischemic stroke.
 - Inhibition of apoptotic cell death**
 - Apoptosis may contribute to a significant proportion of neuron death following acute brain ischemia




Pacific University
Oregon



82

Acute Stroke Management

- N=152
- Open-label, evaluator masked study
- Minocycline 200 mg QD x 5 d or placebo
- Evaluated on NIH Stroke Scale
 - 0-1 complete/nearly complete improvement
 - 2-7 – mild
 - 8-14 – moderate
 - >15 – severe
 - Day 30: 1.8 versus 7.1



Total NIH Stroke Scale Score

1a - Level of Consciousness:	1
1b - LOC Questions:	1
1c - LOC Commands:	1
2 - Best Gaze:	0
3 - Visual Fields:	0
4 - Facial Palsy:	2
5a - Left Motor Arm:	2
5b - Right Motor Arm:	0
5c - Left Motor Leg:	1
5d - Right Motor Leg:	0
7 - Limb Ataxia:	0
8 - Sensory:	1
9 - Best Language:	0
10 - Dysarthria:	1
11 - Extinction and Inattention:	0

Total NIHSS Score: 10

Home Reset All

Lamp I, Boaz M, Gilad R, Lorberboym M, Dabby R, Rapoport A, et al. Minocycline treatment in acute stroke. *Neurology*. 2007;69(14):1404–10

86

TEST	Admission	Day 7	Day 30	Day 90
NHSS - Min	7.5	6.5	1.8	1.6
NHSS - Cont	7.6	8.1	7.3	6.5
mRS - Min	2.8	1.5	1.1	0.9
mRS - Cont	2.0	3.1	2.7	2.1
BI - Min	70.0	85.9	90.6	94.9
BI - Cont	63.9	61.9	68.5	77.6


Minocycline for acute stroke treatment: a systematic review and meta-analysis of randomized clinical trials. *J Neurol*. 2018 Aug;265(8):1871-1879

83

84

Case Report

- 77 YOM
- Right occipital infarct
- 3 weeks post stroke
 - Minocycline 100 mg BID x 5 days




Mark Tomsik, OD and Marlene Skulskie, OD

Pacific University

85

Shortly after TX



1 Year Later



Pacific University

86

Tetracyclines

- This group includes:
 - Tetracycline (250mg - 500 mg cap BID-QID) needs to be taken 1 hour before or 2 hours after a meal.
 - Minocycline (100 mg cap BID)
 - Doxycycline (20mg - 100 mg cap or tab BID)
 - In Canada: Aprillon (30 mg doxy + 10 mg slow release doxy)
- Rules of Thumb with Doxy:
 - Do not take before lying down (>2 hours before)
 - Do not take with calcium and avoid antacids
 - Do not take with dairy
 - Do take with food
 - Do educate on sun protection



87

Side Effects of Tetracyclines

- Side effects include gastric discomfort, phototoxicity, effects on calcified tissues, vestibular problems, IIH.
- Pregnancy Category D.
 - Tetracyclines are attracted to embryonic and growing bone tissue.
 - Depress growth of long bones in pregnant women/children.
 - Cause changes in both deciduous and permanent teeth during the time of tooth development (includes discoloration and increased cavities)
- Contraindicated in:
 - Women in the last half of pregnancy
 - Lactating women
 - Children under 8 years of age



88

Tetracyclines

- Drug of choice for Rocky Mountain Spotted Fever, Cholera, Lyme disease, mycoplasma pneumonia, and chlamydial infections
 - Rocky mountain spotted fever: tick born disease that can potentially be fatal. Doxycycline is drug of choice.
 - 2021 CDC STI Guidelines: doxycycline 100 mg BID for 7 days is the new recommended treatment for chlamydia over azithromycin 1 gram.
- Side effects include gastric discomfort, effects on calcified tissues, vestibular problems.



89

Tetracyclines

- Traditional wisdom is that all tetracyclines should not be used in children under the age of 8 due to discoloration of teeth.
 - Six studies assessed tooth discoloration in at least 338 patients exposed to doxycycline before 8 years of age.
 - Six patients had potential discoloration, but studies consistently found no difference in tooth discoloration between exposed patients and controls.
 - Recommendations have changed stating doxycycline, but not other tetracyclines, can be used for short courses (<21 days) regardless of age.
 - Clinicians should be aware of these data because doxycycline use may extend to disease states apart from tick-borne illnesses in pediatric patients.
- Stultz JS, Elland LS. Doxycycline and Tooth Discoloration in Children: Changing of Recommendations Based on Evidence of Safety. *Annals of Pharmacotherapy*. 2019;53(11):1162-1166.



90

Tetracyclines

- MRSA: methicillin resistant staph aureus
 - Infections with MRSA occur in three specific groups of people: (1) persons currently in the hospital (hospital-associated MRSA), (2) persons with recent hospitalization or ongoing contact with medical clinics, dialysis units, or those undergoing complex outpatient treatments, such as chemotherapy (health care-associated MRSA), and (3) persons in the community (community-associated MRSA).
 - Treatment of MRSA at home usually includes a 7- to 10-day course of an antibiotic (by mouth) such as trimethoprim-sulfamethoxazole (brand name: Bactrim), clindamycin, minocycline, linezolid, or doxycycline.
 - Minocycline: 200 mg orally once, then 100 mg orally twice daily
 - Doxycycline: 100 mg orally twice daily



91

Meibomian Gland Dysfunction

- Meibomian gland dysfunction:
 - also referred to as meibomitis and patients experience dry eye problems secondary to increased evaporation of the tears.
 - signs include noticeable capping of the glands and frothing of tear film.
- Standard treatment includes:
 - good lid hygiene with warm compresses and lid scrubs in conjunction with
 - doxycycline 50 mg po BID for 2-3 months
 - Alternative treatment:
 - Azithromycin 500 mg/day for 3 days for three- four weeks
 - Recent study used single Z-pak treatment



92

Acne Rosacea

- Acne rosacea:
 - affects females > males after 30 with peak incidence 4-7th decade of Celtic/Northern European descent. Males more disfigured.
- 4 subtypes with classic signs of flushing, papules or pustules usually in crops, telangiectasia.
 - **secondary ocular complications (85% of patients)** and often precede other skin manifestations include erythema, itching and burning.
 - Lipases secreted by bacteria on the skin metabolize sebum and produce metabolites that result in inflammation of the skin



93

Acne Rosacea and Demodex

- Demodex is a natural part of human microbiome
- *Demodex folliculorum* live in hair follicles, primarily on the face, as well as in the meibomian glands of the eyelids;
- *Demodex brevis* live in the sebaceous glands of the skin.

94

Acne Rosacea and Demodex

- *Demodex folliculorum* frequently occur in greater numbers in those with rosacea and this overabundance is thought to trigger an immune response or possibly certain bacteria associated with the Demodex

95

Acne Rosacea Management

- Traditionally, treatment for acne/ocular rosacea has focused on symptom suppression to improve patient quality of life and to help manage the disease.
- **Treatment should be initiated even in patients with mild disease, as early intervention can be key in minimizing both the progression of rosacea and its effects on symptoms and visual function.**
- Nonpharmacologic interventions may be useful for the management of the cutaneous manifestations of rosacea.
 - **Avoidance of triggers including gentle skin care, sun-protection and avoiding the use of cosmetic products known to cause flushing.**
 - Treatment options for ocular rosacea include omega-3 fatty acids (FAs) and gamma linolenic acid (GLA), lipid-based artificial tears, lid hygiene with a mild cleanser, hypochlorous acid, and warm compresses.
 - Omega-3 FAs and GLA supplements have been shown to reduce symptoms, lid margin inflammation, and meibomian gland dysfunction (MGD).
 - **Hypochlorous acid predicts ocular hyperinflammation and anti-oxidant properties that decrease the signs and symptoms of ocular irritation that occur when there is an excessive number of bacteria on the lid.**
 - use a tea tree oil-based soap to wash the entire face in order to get the Demodex under control.



96

Acne Rosacea Management

- **Tetracycline, doxycycline, and minocycline have been used for many years for the management of rosacea.** These agents are most useful for improving inflammatory papules and pustules, and may also reduce erythema
 - Since no definitive microbial cause of rosacea has been identified, the efficacy of oral antibiotics in rosacea is often attributed to their anti-inflammatory properties
 - **Support for the efficacy of subantimicrobial doses of doxycycline (20 mg taken twice daily or a combination pill containing 30 mg of immediate release doxycycline and 10 mg of delayed release doxycycline taken once daily) come from both randomized trials and an open label study of almost 1200 patients**



97

Acne Rosacea Management

- **The pharmacologic agent with the strongest evidence for efficacy for persistent facial erythema in rosacea is topical brimonidine (brimonidine tartrate gel 0.33% (brand name: Mirvaso).**
 - Effects can be seen as quickly as 30 minutes after application
 - MIRVASO topical gel is for topical use only and not for oral, ophthalmic, or intravaginal use.
 - **NOT CHEAP: Good Rx Price: \$38 (full)**
- Topical metronidazole, azelaic acid, and topical ivermectin are also considered first-line therapies in mild to moderate facial disease
- **Laser and light-based therapies, which have been used extensively for the treatment of a variety of vascular lesions, have also been used for the vascular features of rosacea, especially telangiectasia.**
 - The mechanism of actions are well known and include immunomodulation of the inflammatory burden, destruction of Demodex, decrease of the bacterial load, photomodulation of mitochondrial activity and rejuvenation of collagen production.



98

Herpes Zoster

1. **Primary infection – Chicken pox (Varicella)**
 - Usually in children
 - Highly contagious***
 - Very itchy maculopapular rash with vesicles that crust over after ≈ 5 days
 - **96% of people develop by 20 years of age**
 - Vaccine now available



99



Herpes Zoster

Reactivation – Shingles (Herpes Zoster)


- More often in the elderly and immunosuppressed (AIDS)
- Systemic work-up if Zoster in someone < 40
- Can get shingles anywhere on the body
- Herpes Zoster Ophthalmicus (HZO)
- Shingles involving the dermatome supplied by the ophthalmic division of the CNV (trigeminal)
 - 15% of zoster cases




100

Herpes Zoster

- Associated factors include increasing age, immune deficiency and stress.
- Traditionally thought to only affect patients over the age of 60 and those patients under 60 should be worked up for immune deficiency
- **Increasing trend to affect patients of younger age who are not immunocompromised**



Shingles and Postherpetic Neuralgia Rates* by Age, United States



Shingles Rates in Adults 30 and Older, 1998-2016

101

Herpes Zoster

- Symptoms:
 - Generalized malaise, tiredness, fever
 - **Headache, tenderness, paresthesias (tingling), and pain on one side of the scalp**
 - Will often precede rash
 - Rash on one side of the forehead
 - Red eye
 - Eye pain & light sensitivity



102

Herpes Zoster


- Signs:
 - **Maculopapular rash -> vesicles -> pustules -> crusting on the forehead**
 - **Respects the midline*****
 - Hutchinson sign
 - **Rash on the tip or side of the nose*****
 - Classically does not involve the lower lid
 - Numerous other ocular signs



103

Herpes Zoster

- Other Eye Complications (Acute):
 - **Anterior uveitis (most common ocular manifestation)**
 - Acute epithelial keratitis (pseudodendrites)
 - Conjunctivitis
 - Stromal (interstitial) interstitial keratitis
 - Endotheliitis (disciform keratitis)
 - Neurotrophic keratitis



104

Herpes Zoster

- Associated factors include increasing age, immune deficiency and stress.
- Only people who had natural infection with wild-type VZV or had varicella vaccination can develop herpes zoster.
- Children who get the varicella vaccine appear to have a lower risk of herpes zoster compared with people who were infected with wild-type VZV.



105

Herpes Zoster

- A person's risk for herpes zoster increases sharply after 50 years of age.
- Almost 1 out of 3 people in the United States will develop herpes zoster during their lifetime.
- A person's risk of developing post-herpetic neuralgia also increases sharply with age.



106

Herpes Zoster

- Management includes:
 - oral antivirals:
 - 800mg acyclovir 5x/day
 - valacyclovir (Valtrex) 1g TID,
 - famciclovir (Famvir) 500 mg TID
 - effectiveness of therapy is best started within 72 hours
 - oral steroids (clinical trials show variable results but often prescribed with antiviral to reduce pain)
 - management of pain (capsaicin, tricyclic antidepressants, gabapentin).
 - If ocular complications, consider topical steroids (Pred Forte QID).



107

NEW!! Shingrix HZ Vaccine

- Approved in US/Canada as of October 2017
- non-live antigen, to trigger a targeted immune response, with a specifically designed adjuvant to enhance this response and help address the natural age-related decline of the immune system
- Shingrix is 97% effective against shingles for people between the ages of 50 and 69 and 91% effective for people 70 or older
- It is 91% effective against postherpetic neuralgia for people 50 and older
- These rates are based on evidence presented to the committee from clinical trials with over 38,000 total participants.



108

NEW!! Shingrix HZ Vaccine

- recommended for healthy adults aged 50 years and older to prevent shingles and related complications
- recommended for adults who previously received the current shingles vaccine (Zostavax®) to prevent shingles and related complications
- the preferred vaccine for preventing shingles and related complications



109