## The Greatest Anterior Segment Disease and Contact Lens Complications Course Ever

Marc Bloomenstein OD FAAO Paul Karpecki OD FAAO Jack Schaeffer OD FAAO Laura Periman MD

1

5

3

Laura Periman MD disclosures FINANCIAL DISCLOSURE FORM
DR JACK L. SCHAEFFER

I HAVE RECEIVED HONORARIUM, COMPENSATION, OR SERVE AS AN ADVISOR TO THE
FOLLOWING COMPANIES

• ALCON
• ALLERGAN
• AMO/ABBOTT
• ARCTIC/DX
• ATON
• BAUSCH AND LOMB
• COOPERVISION
• ESSILOR
• ISTA
• HOYA
• OPTOVUE
• OPTOS

VISTAKON

ZEIS VISION

2

6

#### Dilation Vs Optomap

- The two together delivers a the highest level of Comprehensive Eye Care
- If you have to choose just one:
   DILATE, DILATE, DILATE

#### **Telephone Consultations**

30 YO WF

Telephone symptoms: sore upper lid, painful spot on lid

Internal Hordeolum??

Ready to Dx on telephone: decided to see the patient

7

#### **Bacterial Conjunctivitis?**

Extremely Tender Upper lid
Upper lid swelling
Excessive Mucous production

## Bacterial Conjunctivitis Orbital Cellulitis?

Tx: PO

Augmentin PO 875 Mg Bid

Ocular Zymaxid OS q 2 h

9

#### Day 2

Facial Pain Headache
Fever
Referral to PCP, R/O Orbital Cellulitis
Dx Severe Sinus infection:
Contd Meds PO ( Augmentin)
Antibiotic Injection in office
Sinus infection
Lid swelling with Pain

- Admitted for pain control and IV antihiotics
- Proptosis, edema and pain with eye movement progressed despite broad spectrum IV antibiotics
- · Urgent DCR performed

8

10

11 12

## Chronic Unilateral Conjunctivitis

- 63 yo male with rosacea on chronic doxycycline referred with 'recurrent eye infontional'
- Topical antibiotics would clear symptoms 'a little'
- · Cultures grew out candida species
- Altered ENT microflora with chronic downstline



13 14

Lid Disease-Infection

**Treatment** 

**Chronic Chemosis** 

after Blepharoplasty

- · Keflex 500 Mg BID
  - Cephalexin
- Bactrim: double strength: BID
   Trimethoprim/ Sulfamethoxazol
- Augmetin 875 mg BID
- Miboflow

16

· Hot compress (Written instructions)

15

#### Treatment

- Keflex 500 Mg BID
  - Cephalexin
- Bactrim: double strength: BID
  - Trimethoprim/ Sulfamethoxazol
- Augmetin 875 mg BID
- Miboflow
- · Hot compress (Written instructions)

Chalazia

# Periman IPL Protocol Prep: Trader Joe's or Simple micellar make up remover wipes. Commercial eyelid wipes as needed. Proparaciane, PF AFT, sterifized laser-grade corneal shalds, thin-medium layer of clear ultrasound gel (take great care to avoid gel getting into eyes), applied with long adge for forague depressor. Step 1: Full lace rosaces pass (choose either telangiectasis or erythema based on clinical findings). Step 2: Toyos settings tragus to tragus, double pass Step 3: Switch to small light guide, treat lids, avoid eyelsahes by 2mm, double pass Step 3: Switch to small light guide, treat lids, avoid eyelsahes by 2mm, double pass Step 4: Aesthetic clean up: angioma (VL presets), facial telangiectasis (VL presets), challatis etc. For challatis, stex 4 extra Toyos publes. Post-Procedure: remove gel with ling edge of tongue degressor, gauze remove residual (again, take great care no gel gets into eyes), wipe with warm water. Pat into skin one drop Alphagan P mixed with EttaMD or Skin Medica tinted sunscreen. Place 1:16 dilution of Alphagan P in Refresh Mega into eyes.

#### Caniliculitis/Dacryocystitis

19

#### **Treatment**

- Keflex 500 Mg BID
  - Cephalexin
- Bactrim: double strength: BID
   Trimethoprim/ Sulfamethoxazol
- Augmetin 875 mg BID
- · Hot compress (Written instructions)
- MiBo Flow

21 22

#### **Concretions Management**

- Asymptomatic- neglect (@ 6% become symptomatic
- Symptomatic
  - Fine tipped forceps delivery
  - 25 ga needle
  - Education R.E. recurrence

#### Doctor number 3

• 68 YO female

20

- Treatment
- Pain discomfort 2 years OU
- Restasis BID
- OD > OS
- 3 rd doctor

#### **Allergic Dermatitis**

- Elocon
- · Mometasone Crème
- · Lotemax ung

# Rosacea

**Eyelash Complications** 

25 26

## Hygiene suffers Patients don't want to rub off their expensive

extensions

Prostaglandin Analogs in OTC Eyelash Serums

- Aerodynamic compromise
- Upper eyelid discoloration
- MGD\*
- Hyperpigmentation
- Dermatitis
- Orbital Fat Atrophy
- Iris Color Change
- CME

28

27

MG toxicity from ink pigments	"I'll Look Great Later and
-Tattoo inks may be made from titanium dioxide, lead, chromium, nickel, iron oxides, ash, carbon black, and other ingredients. Some of the pigments are industrial grade and used as automobile paint	No Harm Done"
MG trauma	
Loss of lid margin architecture	Laura M Periman MD
MG dropout	<u> </u>
high SPEED scores	

Trauma / Abrasion

#### **Corneal Abrasion**

- · Debridement of the Cornea
- Techniques
- Instruments
- Bandage Contact lenses
- · Follow up prtocols

31

#### Follow up protocols

- Day 1
- Day 3
- If any symptoms post day 3
  - Telephone
  - Office visit

33

- · Recurrent erosion
  - Any preventive measures

Bandage contact lens

- Pros
- Cons
- Cyclo
- Antibiotic
- Nsaids

32

- Excessive pain
  - Treatment
  - Bandage
  - Nsaid
  - Narcotics
  - Cycloplegia

34

36

**Recurrent Erosion** 

#### **Recurrent Erosion**

#### EBMD / ABMD

#### Case 2

- 50 YOF
- · Woke up with discomfort
- · Feels like something is in my eye

37 38

#### Case 3

- 50 YO male
- Punched in eye 3 weeks ago
- · Ocular Contusion with no abrasions
- Va 20/40

**Treatment Strategy** 

- ABMD
- RCE

40

39

#### **Recurrent Corneal Erosion**

- NaCl Ung Pm
  - Muro 128
- PF AT
  - Q 1-2 hours
- · NaCl Gtts qid

Recurrent Corneal Erosion Long Term Therapy

- Restasis / Xiidra
  - Tid
- · Fresh -Kote
  - Qid
- · Lacriserts?
- Hypertonic Vs Hypotonic AT

- · Amniotic membrane
  - Corneal specialists not in favor
  - Any EBM to support
- · Which type
  - How long to remain in eye
  - Follow up protocol

43

### Recurrent Corneal Erosion

- Bandage Contact lenses
- · Antibiotic ung
- · Change lens how often
- · See patient how often

45 46

#### Epidemic Keratoconjunctivitis

#### Treatment

- · Nsaid?
  - Delayed Corneal healing?
- · Bandage Contact lens
  - Antibiotic??
  - How often
  - RTC daily until healed? How often?
  - Remove and fresh lens and leave in place 3 days?

44

Developing a Specialty Practice

Cornea Disease

**RPS Adeno Detector** 




47 48

### **Treatment EKC**

- 1 lubricants
- 2 combo antimicrobial / steriod
- · 3 Steroid
- 4 Betadine
- 5 Zirgan
- · Contagious ? How long

**EKC** treatment Melton/Thomas · Povidone- Iodine 5% (betadine) - Broad spectrum microbiocide Indicated for "Irrigation of the ocular surface" - OFF LABEL USE · Anesthetize with proparacaine · Instill 1-2 drops NSAID . Instill several drops of betadine in eye ( close eye) Swap excess over lid margin · After one minute irrigate with saline Instill 1-2 drops NSAID Rx Lotemax or Zylet or Tobadex ST qid 4 days - No reports of adverse reactions Avoid if allergic to iodine - Betadine 5% ophthalmic prep soln ( 30 ml opaque) - 99070 supply code


49 50



**Treatment** Zylet qid OD RTC 1 day Some improvement over the new few days, but minimal. SCHAEFFER EYE CENTER

51 52

Treatment D/c Zylet qid OD, begin Besivance q1h OD Differentials? Pt showed significant improvement, at 1-day follow up




53 54



#### **THYGESSONS**

55 56



Thygeson's SPK
 Described by Phillips Thygeson in 1950
 Slightly elevated corneal lesions, minimal staining
 Usually bilateral, Second to third decade
 Noted corneal sensitivity decreased but not as severe as herpes
 Mild conjunctival involvement, worse with exacerbations
 Appearance similar to EKC described by Fuchs

57 58

Thygeson's SPK

• Lesions in basal epithelial layer /
Bowman's layer


• Debris from necrosis / degenerated epi
cells

• Increased Langerhans cell density

• Part of inflammatory response- Type II

Thygeson's SPK Treatment: Anecdotal
 Cyclosporin 2% in olive oil (8 patients)
 Supratarasal injection triamcinolone (1 case-chronic 6+ years)
 Trifluridine (6 eyes)
 PRK in myopic patient had lesions recur in periphery (untreated area) vs central (treated area)
 Rimexolone 1% for reversing dendritic cell density (4 patients)

59 60



Back to the case...
D/c All meds
Lesions healed in 1 week
No recurrences since October

61 62

#### Plaquenil Keratopathy

## Vortex Keratopathy or Cornea Verticillata Clinical features: Symptoms: the corneal changes are rarely of any visual significance. Signs: Symmetric, bilateral, whorl-like pattern of powdery, white, yellow or brown corneal epithelial deposits Appears in a vortex fashion in the inferocentral cornea and swirls outwards sparing the limbus Cocurs in Fabry's disease and in patients being treated with a variety of drugs including amiodarone, chloroquine, amodiaquine, meperidine, indomethacin, chlorpromazine and tamoxifen.

63 64

Ocular Surface Disease Secondary to Systemic Disease

#### Herpes Zoster Management

- Oral antiviral agent
  - Zovirax (acyclovir) 800 mg 5x / day x 7-10 days
  - Famvir (famciclovir) 500 mg tid x 7-10 days
  - Valtrex (valacylovir) 1000 mg tid x 7-10 days
  - Discussed with nephrologist / PCP if renal disease present

65 66

#### **Ocular findings:**

- Conjunctivitis/Scleritis
- Pseudodendrites
- Neurotrophic keratitis
- Iritis
- Glaucoma
- ION, vein or artery occlusion
- Nerve Palsy

#### **Iridocyclitis and HZO**

- Most common and most often overlooked ocular complication (43%)
- Highly elevated IOP
- Study by Thean, Hall & Stawall -clinical Ophthalmology Dec 2001
- •56% of patients developed glaucoma!!

67

#### **Treatment:**

- •Duration?
- 7 days for most patients although newer studies (Zaal - Am J or Ophthal. Jan 2001) suggest
- •10 days for patients over age 66 due to shedding

68

70

#### **Treatment: Iridocyclitis**

- •Pred Acetate 1% q1h or q2h or
- •Durezol (Difluprednate) 0.05% with half the dosing
- Lotemax Long term
- Cycloplegia
- Homatropine 5% bid
- Cyclopentolate 1% bid

69

#### Systemic Disease- Ocular Involvement

Herpes Simplex

### Treatment: Epithelial Involvement

- •In the past: trifluoridine Viroptic q2h
- New replacement: Zirgan 5 x per day until ulcer disappears then TID x 1 week
- PO Valtrex 500mg TID
- PF artificial tears
- •Follow-up (next day), day 3-4, day 7-10

71 72

### Zirgan™(Ganciclovir Ophthalmic Gel) 0.15%

### Zirgan™ (ganciclovir ophthalmic gel) 0.15% Indication

Dosage and Administration

 The recommended dosing regimen for Zirgan is 1 drop in the affected eye 5 times per day (approximately every 3 hours while awake) until the corneal ulcer heals, and then 1 drop 3 times per day for 7 days.

**Restoring Corneal Clarity** 

\* Patient continued oral Valtrex 500mg QD

\* Returns 5 Days later for removal:

\* Prokera inserted

73 74

#### Treatment: Stromal keratitis or Endotheliitis

Durezol QID

75

- Pred Forte Q2H
- Cover with PO Acyclovir (400 mg bid) or Valtrex (1000mg QD) or topical (Zirgan TID)

Inflammation is the Hallmark of All Ocular Surface Diseases

Ocular Surface Disease

Corneal Conjunctival Inflammation Eyelid Inflammation

Keratitis Conjunctivitis Blepharitis

## The Structure of the Fetal Amniotic Membrane • Metabolically active cuboidal cells with microvilli present on its apical surfacce • Made up type IV, V and VII collagen (also found on conjunctival and corneal basement membranes) • Fibronectin and Laminin • Compact Layer provides tensile strength • Fibroblast Layer • Spongy Layer

77 78

13