


Common Ocular Manifestations From Systemic Associations: Front to Back

Mark Schaeffer, OD FAAO




1

Financial Disclosures - Mark Schaeffer, OD FAAO

I Have Received Honoraria From:

- AesculaTech - Consultant
- Alcon - Consultant, Speaker
- Allergan - Consultant, Speaker
- Bausch + Lomb - Consultant, Speaker
- CooperVision - Consultant
- Hermal - Consultant, Speaker
- Johnson & Johnson Vision Care - Consultant
- LENZ Therapeutics - Consultant
- Optase - Consultant
- Science Based Health - Consultant
- Sight Sciences - Consultant
- Tarsis - Consultant
- Zeiss - Consultant
 - Founder, Dr. MES Consulting
 - Founding Member, Intrepid Eye Society



3

How do you manage systemic disease patients?

4

Case #1

5

Comprehensive Eye Exam

39 year old female

- Comprehensive Eye Exam, updating glasses and contact lenses
- During case history, lenses have been bothering her off and on
- Getting more dryness as the end of the day
- Has been to multiple ODs, has been in daily disposables
- Here today for updated Rx
- Medical history unremarkable
- SPEED score- 8


6

Eye Exam and findings

	OD	OS
Refraction	Plano -2.25x015 20/15	-0.50 -1.00x176 20/15
Slit lamp exam	Decreased tear break up time (+) NaFL staining inferior cornea	Decreased tear break up time (+) NaFL staining inferior cornea
Cup-to-disc ratio	0.3/0.3	0.3/0.3
Fundus evaluation	(-) H, T, D 360 OU	(-) H, T, D 360

7

Let's look at the cornea...



8

Plan of Action

-Sidebar discussion: Do you discontinue contact lenses?

- Refit into premium Silicone Hydrogel daily disposable contact lens
- Started on Lifitegrast bid OU
- RTC 2-3 months

9

Patient returns to clinic

12 months later

- Patient felt better on medication
- Contact lenses felt better at the end of the day
 - Patient commented, "This is what usually happens for a little bit"
- Visual acuity 20/15 OD, OS, OU
- Patient felt better after 2-3 months, discontinued the treatment and is now feeling worse

10

Do we want to ask some more questions?

11

Systemic questions

- Do you suffer from dry mouth?
- Any other areas of dryness?
- Do you frequently have to drink liquids when swallowing food?

12

Patient responses

- Points to water bottle
 - "I can't go anywhere without it. Constantly consuming water"
- Was planning on discussing with OB/GYN regarding vaginal dryness

13

Patient encouraged to get testing from PCP

14

Current plan


- Patient resumed Liftegrast
- On shorter return schedule for compliance
- Returns in a couple months
 - "Did you schedule with your primary care?"
 - (No response)
- "Do you want us to schedule with your primary care?"
- (No response)

15

COVID happens

16

Then I get this in the mail to my office...




Dr. Schaeffer,
I wanted to write to tell you how often I feel grateful for your care and recommendation. I did not bleed well and was diagnosed with Sjogren's syndrome. I completely attribute an early diagnosis to your persistence of me getting blood work done. THANK YOU! I had your fortunate that it only affected my eyes prior to starting treatment which should slow and hopefully stop progression. Again thank you for being great at what you do!

17

Sjogren's Syndrome

- Auto-immune deficiency of body's salivary glands to produce fluid
- Of individuals with significant aqueous deficient dry eye, 10% are likely to have Sjogren's syndrome
- 26% of patients with either aqueous tear deficiency or evaporative dry eye have an underlying rheumatic condition, including Sjogren's syndrome
- Of 1208 participants in international Sjogren's syndrome registry, 85% reported symptoms of dry eye



18

New questions in Sjogren's

1. Is your mouth dry while eating a meal?
2. Can you eat a cracker without drinking any fluids?
3. How often do you have excessive tearing?
4. Are you able to produce tears?

19

One more on the “surface”

20




21

Thyroid Eye Disease

- Auto-immune disease of the eye and orbital muscles
- Characterized by acute, progressive inflammatory stage then chronic, fibrotic stage 1-3 years later
- Typically found in patients with Grave's disease (40% have TED)
- Can be found in patients with hypothyroid and euthyroid as well

22



23

Signs of TED

- Orbital congestion (not to be mistaken for conjunctivitis)
- Allergic conjunctivitis without any papillary reaction that doesn't improve with allergy drops
- Unexplained changes in vision that are inconsistent with corneal changes from dryness or other pathologies, which can actually be caused by low-grade chronic compressive optic neuropathy
 - Resistance to retropulsion, an unsatisfactory response to a careful motility check and lid lag on down-gaze can help with this diagnosis
- Optic nerve imaging with OCT and visual field testing can be helpful in these cases

24

TED is a Condition of the Eye, But Also the Face

In addition to orbital soft tissue expansion, TED is associated with expansion of soft tissue in other regions of the face, including the brow, temples, and mid and lower face

Tepratumumab reduced mean soft tissue volume across facial regions (n=23 patients with mean TED duration of 29 months)


59 year-old female with 10-month history of TED and expansion of facial soft tissue

25


Clinical Experience with Teprotumumab for Chronic TED

- Patient selection
 - 68-year-old Black male with a 9-year history of TED
 - Previous treatment with steroids, bilateral orbital radiation, bilateral orbital decompression, and strabismus surgery
 - Progressively worsening proptosis
 - Exophthalmometry: 32 mm OD, 32.5 mm OS
 - CAS of 3
- After 8 infusions of teprotumumab, the patient saw clinically significant reductions in proptosis and reached a CAS of 0
 - Exophthalmometry: 26 mm OD, 28 mm OS
 - Reduction of 6 mm and 4.5 mm, respectively

Before teprotumumab treatment



After teprotumumab treatment



OD, right eye; OS, left eye; CAS, Clinical Activity Score

Images shown in published [1] and formed under a Creative Commons Attribution 4.0 International License

26

Cell-ebrate Good Times!



27

27 year old female reports to the clinic

Red Eye / Urgent care appointment

- Complains of pain, redness, and light sensitivity in the right eye
- Started yesterday, but has gotten worse
- Used Visine but didn't help
- Medical history unremarkable*
- Denied prescription medication use, but uses NSAID for headaches, muscle pain

28

Representative Photos

29

Uveitis

Anterior Granulomatous Non-infectious Uveitis OD

- Patient was started on difluprednate qid OD
- Dosed with Cyclopentolate 1% in office
- RTC 24 hours
- Referral to PCP for lab work


- Resolved with steroid treatment
- Quick aside on tapering with a steroid

30


Patient (+) RF, ANA Working diagnosis: Systemic Lupus Erythematosus

31


LAB WORKUPS



WHEN DO YOU ORDER LABS?



WHAT DO YOU TELL THE PATIENT?



HOW DO YOU CO-MANAGE WITH SUB-SPECIALTIES?

32

WHEN SHOULD LAB TESTS BE ORDERED?

- Bilateral cases
- Hyperacute cases
- Atypical age group
- Worsens with tapering
- Recurrent uveitis
- VA worsening
- Recalcitrant cases
- Immunosuppressed px

33

WHAT LABS TO ORDER?

- *Complete Blood Count with differential (CBC with diff)
- *Erythrocyte Sedimentation Rate (ESR)
- *C Reactive Protein (CRP)
- *Anti-Nuclear Antibody (ANA)
- *Angiotensin Converting Enzyme (ACE)
- Chest X-ray (CXR)
- Anti-neutrophil Cytoplasmic Antibody (ANCA)
- *Venereal Disease Research Laboratory (VDRL) / Rapid Plasma Reagin (RPR)
- *Fluorescent Treponema Antibody Absorption Test (FTA-Abs)
- *Purified Protein Derivative (PPD) or Mantoux Test
- Enzyme Linked Immunosorbent Assay (ELISA) or Immunofluorescence Assay (IFA)
- Western blot
- *Rheumatoid Factor (RF)
- *Human Leukocyte Antigen (HLA)
- CD4 count

34

SYSTEMIC LUPUS ERYTHEMATOSUS

Chronic, microvascular inflammatory condition

- Generation of autoantibodies
- Defect in self-apoptosis signal
- Increased cell death

Epidemiology

- 6.1 per 10,000 (number on the rise)
- Average 6 years from first symptoms to diagnosis

Patient Profile

- F>>>M (2:1 and 11:1 during child-bearing years)
- Affects minorities more


Prognosis

- Had 50% mortality rate previous, has decreased to 10%

Diagnostic Testing

- Anti-Nuclear Antibody (ANA)

35



36

Case


- Dry Eye Consult
- 35yo F
- CC; Blurred vision

37


Case

- CC; Blurred vision
- Lupus, Sjogrens
- "Sparkly" Vision
- Kidney issues


38



39



40



41

Chloroquine and Hydroxychloroquine Retinopathy

Recommendations on Screening for Chloroquine and Hydroxychloroquine Retinopathy (2016 Revision)

Michael F Marmor¹, Ulrich Kelliner², Timothy Y Y Lai³, Ronald B Melles⁴, William F Mieler⁵; American Academy of Ophthalmology

42

Chloroquine and Hydroxychloroquine Retinopathy

- Asian patients often show an extramacular pattern of damage.
- DOSE: We recommend a maximum daily HCQ use of ≤ 5.0 mg/kg real weight, which correlates better with risk than ideal weight.
- There are no similar demographic data for CQ, but dose comparisons in older literature suggest using ≤ 2.3 mg/kg real weight.

43

Chloroquine and Hydroxychloroquine Retinopathy

- The risk of toxicity is dependent on daily dose and duration of use. At recommended doses,
- The risk of toxicity up to 5 years is under 1%
- And up to 10 years is under 2%,
- But it rises to almost 20% after 20 years.
- However, even after 20 years, a patient without toxicity has only a 4% risk of converting in the subsequent year.

44

Chloroquine and Hydroxychloroquine Retinopathy

- A baseline fundus examination should be performed to rule out pre-existing maculopathy.
- Begin annual screening after 5 years for patients on acceptable doses and without major risk factors.

45

Chloroquine and Hydroxychloroquine Retinopathy

- High dose and long duration of use are the most significant risks.
- Other major factors are concomitant;
 - Renal disease
 - Tamoxifen.

46

Chloroquine and Hydroxychloroquine Retinopathy


- The primary screening tests are
 - Automated visual fields (10-2 STD)
 - Spectral-domain optical coherence tomography (SD OCT).
 - The multifocal electroretinogram (mfERG) can provide objective corroboration for visual fields
 - Fundus autofluorescence (FAF) can show damage topographically

47

Chloroquine and Hydroxychloroquine Retinopathy

- Retinopathy is not reversible, and there is no present therapy.
- Recognition at an early stage (before any RPE loss) is important to prevent central visual loss.
- However, questionable test results should be repeated or validated with additional procedures to avoid unnecessary cessation of valuable medication.

48



49

53 year old male reports for eye exam


- Complains of blurred vision, worse in the evening in the right eye
- History of Type 2 diabetes, "controlled" on medications
 - A1c: 8.2
 - Taking Metformin, Glipizide
- Previous fundus examinations unremarkable for diabetic retinopathy with similar A1c

50

Eye Exam and findings

	OD	OS
Refraction	-4.25-0.50x084 20/25-	-3.00-0.25x096 20/20+
Slit lamp exam Pre-dilation	Healthy, unremarkable	Healthy, unremarkable
Cup-to-disc ratio	0.25/0.25	0.25/0.25
Fundus evaluation	(+) hemes, exudates, cotton wool spots, NVD, NVE	(+) hemes, exudates, cotton wool spots, NVD, NVE


51



52

Diabetic Posterior Subcapsular cataract

53



54


Why diabetes can affect the lens

- Protein build-up in the lens
- Can change the index of refraction leading to fluctuating prescriptions
 - Bag of glasses

55

Do you test A1c in your office?

56



57

Patient findings

	OD	OS
BCVA	20/15	20/15
IOP	18mmHg	17mmHg
Cornea	Clear	Clear
Conjunctiva	White and quiet	White and quiet
Iris / Ant Chamber	Deep and quiet	Deep and quiet
Lens	Clear	Clear

58


Patient findings continued

	OD	OS
Discs	2/2, large area of PPA	2/2, large area of PPA
Vitreous	Clear	Clear
Macula	Normal	Normal
Retina	See photo	See photo

59



60



61



62



63

Next questions

- No nausea, headaches, dizziness, systemic symptoms
- Feeling good today
- Hasn't noticed any changes in vision or otherwise
- Discontinued lisinopril 3-4 months ago

64

Hypertensive retinopathy

Are you a numbers person or a qualitative staging person?

Grade of retinopathy	Retinal findings	Guidance for therapy	Prognosis
None	generalized arteriolar narrowing or sclerosis	< 130 and < 130	Optimal management with risk reduction
Mild	AV crossing changes microhemorrhages, cotton wool spots, flame hemorrhages, hard exudates, soft exudates, retinal edema, venous beading	< 130/130	Optimal management with risk reduction
Severe	retinal hemorrhages, exudates and cotton wool spots	> 130	Optimal management with therapy
Grade IV	Sclerosis and optic disc edema of retinal arterioles Severe grade III and papilloleptema		

From Shegolev et al.
doi:10.1001/archophth.117.10.1411

65

Grade 3 (Moderate) Hypertensive Retinopathy

66

Next steps

- Go directly to the ER!
- Patient lost to follow-up

67

9 months later...

28 year old new patient reports to the office
"Doc, you don't know me but I know who you are. I want to say thank you"

"Because of you, my grandmother got to hold her great granddaughter who was born 4 months ago"


Grandmother went to the ICU for 7 days for impending stroke

68

THANK YOU!

**Common Ocular Manifestations From Systemic Associations:
Front to Back**

Mark Schaeffer, OD FAAO mark@drmesconsulting.com



69